This is a typical member of the *Cortinicara gibbosa* (Herbst) group, species of which are best separated by their characteristic male aedeagi (see Johnson, 1975, 1977a, 1977b).

Acknowledgements

I am indebted to C. Reid and the late J. Klapperich for material collected in Colombia and China respectively, which is now in the Manchester Museum. Other material was kindly loaned by R.D. Pope, British Museum (Nat.Hist.); R. Baranowski and R. Danielsson, Zoological Museum, University of Lund; C. Besuchet and I. Löbl, Muséum d'Histoire Naturelle, Geneva.

References

- Johnson, C. 1972a. Some members of the genera *Corticarina* Reitter and *Melanoph-thalma* Motschulsky (Col., Lathridiidae) from India and Nepal. *Entomologist* **105**: 97-110.
- —. 1972b. Studies on the genera *Corticarina* Reitter and *Melanophthalma* Motschulsky (Col., Lathridiidae). *Nouv. Rev. Ent.* 2: 185-199.
- —. 1975. *Cortinicara*, a new genus of Corticariinae (Coleoptera: Lathridiidae). *Entomologica scand*. **6**: 283-285.
- —. 1977a. Ergebnisse der Bhutan-Expedition 1972 des Naturhistorischen Museums in Basel, Coleoptera: Fam. Lathridiidae Subfam. Corticariinae. *Entomologica basil.* 2: 329-335.
- —. 1977b. Beiträge zur kenntnis der Fauna Afghanistans, Merophysiidae and Lathridiidae, Coleoptera. *Casopis Morav. Muz.* 62: 123-132.
- —. 1979. New species of *Corticarina* Reitter (Col., Lathridiidae). *Entomologist's mon. Mag.* (1978) 114: 55-62.
- —. 1981. New species of *Corticarina* Reitter (Col., Lathridiidae), Part 2. *Entomogist's mon. Mag.* 117: 117-123.

Brachypterolus linariae (Stephens) (Col.: Nitidulidae) in Surrey

In the past, this species was confused with *B. pulicarius* (L.), long recognised as a British insect. In a paper establishing the identity of *B.linariae* (Stephens) and recording its presence also in Britain, my friend Mr C. Johnson (1967 *Entomologist* 100: 142) listed a few locatities for the species but these did not include Surrey. I should like, accordingly, to record that I collected eight specimens of *Brachypterolus* from flowers of *Linaria vulgaris* Mill. from a site near Cheam, Surrey on 15.viii.89. As it turned out, all were males; four proved to be *B. linariae* and 4 were *B. pulicarius*.

The presence of equal numbers of *linariae* and *pulicarius* in this, albeit small, sample, is in marked contrast to what I have found in other districts. There, *pulicarius* has always greatly outnumbered *linariae*. It could be interesting to see, over the next few years, whether the Cheam site maintains this equality in the two species. — J.A. OWEN, 8 Kingsdown Road, Epsom, Surrey KT17 3PU.